

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 797 978 A2

(12)

EUROPEAN PATENT APPLICATION

17

(43) Date of publication:

01.10.1997 Bulletin 1997/40

(51) Int. Cl.⁶: A61K 7/06

(21) Application number: 97105023.2

(22) Date of filing: 25.03.1997

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE

(30) Priority: 29.03.1996 JP 75903/96

(71) Applicant: Kyowa Hakko Kogyo Co., Ltd.
Chiyoda-ku, Tokyo 100 (JP)

(72) Inventors:

- Takahashi, Tomoya
- Tsuchiura-shi, Ibaraki 300-11 (JP)

• Yokoo, Yoshiharu

Ushiku-shi, Ibaraki 300-12 (JP)

• Kamiya, Toshikazu

Machida-shi, Tokyo 194 (JP)

• Shirai, Akio

Machida-shi-shi, Tokyo 194 (JP)

• Tamaoki, Tatsuya

Machida-shi, Tokyo 194 (JP)

(74) Representative: Casalonga, Axel et al
BUREAU D.A. CASALONGA - JOSSE
Morassistrasse 8
80469 München (DE)

(54) Hair-growing agent comprising a specific inhibitor of protein kinase C

(57) Disclosed is a safe and effective hair-growing
agent comprising a protein kinase C-specific inhibitor.

EP 0 797 978 A2

Description**FIELD OF THE INVENTION**

5 The present invention relates to a hair-growing agent comprising a protein kinase C-specific inhibitor.

BACKGROUND OF THE INVENTION

10 There is disclosed that substances with a protein kinase-inhibiting activity, 3-amino/hydroxy-4-[4-benzoylphenyl carbonylamino/oxy]azepanes stimulate the growth of hair (see EP663393 A1); but said protein kinase inhibitors have a protein kinase A (hereinafter referred to as PKA)-inhibiting activity along with a protein kinase C (hereinafter referred to as PKC)-inhibiting activity.

In hair-follicle organ culture systems, a PKC inhibitor, H-7 [1-(5-Isoquinolinylsulfonyl)-2-methylpiperazine] is known to release the hair growth-retarding activity of a PKC-activating substance, 12-O-tetragalloyl-phorbol-13-acetate (see 15 British Journal of Dermatology, 133, 5, 686-693, 1995). Said H-7 is further known to have a PKA-inhibiting activity along with a PKC-inhibiting activity (see BIO/TECHNOLOGY, 8, 732, 1990).

In hair-follicle organ culture systems, the PKC inhibitor, H-7 releases the hair growth-inhibiting activity of 12-O-tetragalloyl-phorbol-13-acetate, but said PKC inhibitor has not been known to have a hair growth-promoting activity (see British Journal of Dermatology, 133, 5, 686-693, 1995).

20 Protein kinase inhibitors having both a PKC-inhibiting activity and a PKA-inhibiting activity could not always be expected to produce satisfactory hair growth-promoting results. H-7 and 3-amino/hydroxy-4-[4-benzoyl-phenyl carbonylamino/oxy]azepanes could not always be expected to produce satisfactory hair growth-promoting results, because such compounds have both a PKC-inhibiting activity and a PKA-inhibiting activity.

The present inventors have first found that PKC-specific inhibitors produce satisfactory hair growth-promoting 25 results.

SUMMARY OF THE INVENTION

30 The object of the present invention is to provide a safe and effective hair-growing agent comprising, as an active ingredient, a PKC-specific inhibitor, and pharmaceutically acceptable vehicles.

DETAILED DESCRIPTION OF THE PRESENT INVENTION

35 The protein kinase C-specific inhibitor as referred to herein is a protein kinase inhibitor having a PKC-inhibiting activity while having a PKA-inhibiting activity as little as possible. Specifically, any inhibitor can be used so long as the ratio of its 50 % PKA-inhibiting constant (hereinafter referred to as PKA-IC₅₀) to its 50 % PKC-inhibiting constant (hereinafter referred to as PKC-IC₅₀), PKA-IC₅₀/PKC-IC₅₀, is not smaller than 3.0 when determining PKC-IC₅₀ and PKA-IC₅₀ according to the following PKC-inhibiting activity measuring method and according to the following PKA-inhibiting activity measuring method, respectively. For example, usable herein are protein kinase inhibitors having PKA-IC₅₀/PKC-IC₅₀ 40 of from 3 to 10⁹, and preferred are those having PKA-IC₅₀/PKC-IC₅₀ of from 10 to 10⁹.

For determination of PKA-inhibiting activity of protein kinase inhibitors having a low PKA-inhibiting activity, the following method requires a large amount of such a protein kinase inhibitor to be determined therein, in which, therefore, the protein kinase-inhibiting activities of the protein kinase inhibitor could be measured only up to its uppermost soluble concentration in the system. Accordingly, if such a protein kinase inhibitor with low PKA-inhibiting activity is used, the 45 value of PKA-IC₅₀/PKC-IC₅₀ capable of being numerically determined in said method will be about up to 10⁹. The protein kinase inhibitors usable in the present invention have its value of PKA-IC₅₀/PKC-IC₅₀ of not smaller than 3, without being defined by its upper value of PKA-IC₅₀/PKC-IC₅₀(10⁹) capable of numerically determined by said method.

(1) Method for Measuring PKC-inhibiting Activity:

50

To measure the PKC-inhibiting activity of a protein kinase inhibitor, referred to is the Kikkawa et al's method (see Journal of Biological Chemistry, 257, 13341, 1982).

Precisely, 10 µl of a sample to be tested for determining its activity is added to 250 µl of a solution comprising 2.5 µmols of magnesium acetate, 50 µg of histone Type IIS (produced by Sigma Co.), 20 µg of phosphatidylserine, 0.8 µg 55 of diolein, 25 nmols of calcium chloride, 5 µg of a crude enzyme (as partially purified from a rat brain according to the Kikkawa et al's method) and 5 µmols of Tris-HCl buffer (pH 7.5), and incubated therein at 30°C for 3 minutes.

After the completion of the incubation, 1.25 nmols of [γ-³²P]ATP (from 5 x 10³ to 10 x 10³ cpm/nmol) is added to the system, and phosphorylation reaction is carried out at 30°C for 3 minutes, and thereafter the reaction is terminated by adding 25 % trichloroacetic acid thereto.

The resulting reaction mixture is filtrated through a cellulose acetate membrane (pore size: 0.45 μm) (produced by Toyo Filter Co.), and the membrane is washed four times with 5 % trichloroacetic acid, and thereafter the radioactivity of the residue remained on the membrane is measured to be a value of the sample.

On the other hand, the same process as described above is repeated without adding the sample to the system, and the radioactivity is obtained to be a control value.

The molar concentration of the sample having the value of radioactivity which is 50 % of the control value is obtained to be the 50 % PKC-inhibiting constant (PKC-IC₅₀) of the sample.

(2) Method for Measuring PKA-inhibiting Activity:

To measure the PKA-inhibiting activity of a protein kinase inhibitor, referred to is the Kuo et al's method (see Biochemistry, 64, 1349, 1969).

Precisely, 10 μl of a sample to be tested for determining its activity is added to 250 μl of a solution comprising 5 μmol s of Tris-HCl buffer (pH 6.8), 2.5 μmol s of magnesium acetate, 100 μg of histone Type IIS (produced by Sigma Co.), 0.25 nmol of c-AMP and 200 μg of a crude enzyme (as partially purified from a calf heart according to the Kuo et al's method), and incubated therein at 30°C for 3 minutes.

After the completion of the incubation, 1.25 nmol of [g -³²P]ATP (from 5×10^3 to 10×10^3 cpm/nmol) is added to the system, and phosphorylation reaction is carried out at 30°C for 3 minutes, and thereafter the reaction is terminated by adding 25 % trichloroacetic acid thereto.

The resulting reaction mixture is filtrated through a cellulose acetate membrane (pore size: 0.45 μm) (produced by Toyo Filter Co.), and the membrane is washed four times with 5 % trichloroacetic acid, and thereafter the radioactivity of the residue remained on the membrane is measured to be a value of the sample.

On the other hand, the same process as described above is repeated without adding the sample to the system, and the radioactivity is obtained to be a control value.

The molar concentration of the sample having the value of radioactivity which is 50 % of the control value is obtained to be the 50 % PKA-inhibiting constant (PKA-IC₅₀) of the sample.

Specific examples of the protein kinase C-specific inhibitor for use in the invention may include polymyxin B, calphostin C, palmitoyl-DL-carnitine and hexadecylphosphocholine (militefosine, produced by Sigma Co.), and also their pharmaceutically-acceptable salts.

The pharmaceutically-acceptable salts may include, for example, hydrochlorides, hydrobromides, sulfates, nitrates, formates, acetates, benzoates, maleates, fumarates, succinates, tartrates, citrates, oxalates, methanesulfonates, toluenesulfonates, aspartates, and glutamates.

The hair-growing agent of the present invention may be taken any form, provided that it properly contains the protein kinase C-specific inhibitor of the invention. For example, a liquid or solid hair-growing agent comprising the protein kinase C-specific inhibitor of the invention along with pharmaceutically acceptable vehicles is used.

The liquid or solid hair-growing agent may include liquid-type preparations such as hair liquids, hair tonics and hair lotions; and solid-type preparations such as ointments and hair creams. These agents can be prepared by any ordinary methods, while adding the protein kinase inhibitor of the invention to suitable vehicles.

The amount of the protein kinase C-specific inhibitor to be in the hair-growing agent of the present invention greatly varies, depending on the intensity of its inhibiting activity and also on its endermic absorbability to be derived from its physical properties, but may be generally from 10^{-6} to 10 % by weight (hereinafter referred to as %) based on the agent in terms of the content of a single compound of the inhibitor or a mixture of plural compounds thereof.

The preferred vehicles for the liquid-type preparations are those that are generally used in ordinary hair-growing agents, such as pure water, ethanol, polyalcohols, oils and fats. If desired, any additives may be added thereto.

The polyalcohols may include, for example, glycerol, 1,3-butylene glycol, propylene glycol, etc.

The oils and fats may include, for example, wheat germ oil, camellia oil, Jojoba oil, olive oil, squalane, safflower oil, macadamia nut oil, avocado oil, hydrogenated soybean lecithin, etc.

The additives may include, for example, fragrances, surfactants microbicides, etc. If desired, any of antioxidants, hormones, ultraviolet absorbents, anti-inflammatory agents, refrigerants, moisturizers, vitamins, herb extracts tinctures, etc. may be added to the preparations.

As the fragrances, any fragrance usable in ordinary cosmetics, etc are employable herein.

The surfactants may include, for example, polyoxyethylene(60) hardened castor oil, polyoxyethylene(8) oleyl ether, polyoxyethylene(10) oleyl ether, polyoxyethylene(10) monooleate, polyoxyethylene(30) glyceryl monostearate, sorbitan monostearate, polyoxyethylene(30) glyceryl monostearate, polyoxyethylene(20) sorbitan monooleate, sucrose fatty acid esters, hexaglycerin monooleate, hexaglycerin monolaurate, polyoxyethylene reduced lanolin, polyoxyethylene(20) lanolin alcohol, polyoxyethylene(25) glyceryl pyroglutamate isostearate diester, N-acetylglutamine isostearyl ester, etc.

The microbicides may include, for example, trichlorohydroxydiphenyl ether, hinokitiol, triclosan, chlorohexidine gluconate, phenoxyethanol, resorcinol, isopropylmethylphenol, azulene, salicylic acid, zinc pyrithione, benzalkonium chlo-

ride, photoreceptor No. 301, sodium mononitroguaiacol, etc.

The antioxidants may include, for example, butylhydroxyanisole, propyl gallate, and erysorbic acid.

The hormones may include, for example, ethynylestradiol, estrone, estradiol, etc.

The ultraviolet absorbers may include, for example, benzophenones such as dihydroxybenzophenone; as well as
5 melanins, ethyl para-aminobenzoate, 2-ethylhexyl paradimethylaminobenzoate, cinoxate, 2-ethylhexyl paramethoxycin-
namate, 2-(2-hydroxy-5-methylphenyl)benzotriazole, urocanic acid, and fine particles of metal oxides, etc.

The anti-inflammatory agents may include, for example, dipotassium glycyrrhetinate, β -glycyrrhetic acid, allan-
toin, diphenhydramine hydrochloride, guaiazulene, and 1-menthol, etc.

The refrigerants may include, for example, capsicum tincture, 1-menthol, etc.

10 The moisturizers may include, for example, L-pyrrolidonecarboxylic acid, sodium hyaluronate, chondroitin sulfate,
etc.

The vitamins may include, for example, dl- α -tocopherol acetate, dl- α -tocopherol, d- δ -tocopherol, vitamin E, benzyl
nicotinate, nicotinic acid amide, D-pantothenyl alcohol, pantothenyl ethyl ether, biotin, pyridoxine hydrochloride, ribofla-
vin, etc.

15 The herb extracts may include, for example, Swertia herb extract, garlic extract, ginseng extract, aloe extract, cin-
chona extract, etc.

The tinctures may include, for example, capsicum tincture, ginger extract, cantharis tincture, etc.

Where the above-mentioned liquid-type preparations are used as spray, they may be combined with noninflam-
mable gas or the like.

20 The vehicles for the solid-type preparations may include, for example, vaseline, solid paraffin, vegetable oil, mineral
oil, lanolin, wax, and macrogol. To these may be added, if desired, any of the above-mentioned additives; emulsifiers
such as lecithin; and lower alcohols such as ethyl alcohol isopropyl alcohol, etc.

The amount of the hair-growing agent of the present invention to be applied varies, depending on the age, the body
weight and the condition of the case to which it is applied, the curing effect of said agent, the mode of administration,
25 the treating time, etc. For a hair tonic of the hair-growing agent of the present invention, for example, its percutaneous
dose may be from 0.5 to 5 ml/adult, preferably from 1 to 3 ml/adult, and may be applied once or several times a day.

Now, the present invention is described in more detail hereinunder with reference to the following examples, refer-
ence example and test examples.

30 Example 1: Preparation of Hair-growing agent Tonic 1

First, 55 g of ethyl alcohol, 7 g of 1,3-butyleneglycol, 0.5 g of N-acetylglutamine isostearyl ester, and 0.25 g of poly-
oxyethylene(25) glyceryl pyroglutamate isostearate diester were uniformly mixed and stirred to prepare a solution A.

Next, 0.3 g of polymyxin B sulfate (produced by Sigma Co.) and 36.95 g of pure water was uniformly mixed and
35 stirred to prepare a solution B.

The solution B was added to the solution A with stirring, and uniformly mixed to prepare a hair-growing agent tonic
(composition 1).

Example 2: Preparation of Hair-growing agent Tonic 2

40 First, 90 g of ethyl alcohol, 5 g of 1,3-butyleneglycol, 0.5 g of N-acetylglutamine isostearyl ester, 0.25 g of polyox-
yethylene(25) glyceryl pyroglutamate isostearate diester, and 0.03 g of calphostin C (produced by Kyowa Hakko Kogyo
Co., Ltd) were uniformly mixed and stirred to prepare a solution A.

To the solution A was added 4.22 g of pure water with stirring, and uniformly mixed to prepare a hair-growing agent
45 tonic (composition 2).

Example 3: Preparation of Hair-growing agent Tonic 3

50 First, 77 g of ethyl alcohol, 10 g of 1,3-butyleneglycol, 0.5 g of N-acetylglutamine isostearyl ester, 0.25 g of polyox-
yethylene(25) glyceryl pyroglutamate isostearate diester, and 3 g of palmitoyl-DL-carnitine hydrochloride (produced by
Sigma Co.) were uniformly mixed and stirred to prepare a solution A.

To the solution A was added 9.25 g of pure water with stirring, and uniformly mixed to prepare a hair-growing agent
tonic (composition 3).

55 Example 4: Preparation of Hair Growing Tonic 4

First, 70 g of ethyl alcohol, 10 g of 1,3-butyleneglycol, 0.5 g of N-acetylglutamine isostearyl ester, 0.25 g of polyox-
yethylene(25) glyceryl pyroglutamate isostearate diester, and 1 g of hexadecylphosphocholine(produced by Sigma Co.)
were uniformly mixed and stirred to prepare a solution A.

Then, 18.25 g of pure water was added to the solution A with stirring, and uniformly mixed to prepare a hair-growing agent tonic (composition 4).

Reference Example 1: Determination of PKC-IC₅₀ and PKA-IC₅₀

According to the PKC-inhibiting activity measuring method and the PKA-inhibiting activity measuring method mentioned hereinabove, the PKC-inhibiting activity and the PKA-inhibiting activity of polymyxin B sulfate, calphostin C, palmitoyl-DL-carnitine hydrochloride and hexadecylphosphocholine were measured, from which were obtained their PKC-IC₅₀ and PKA-IC₅₀.

The data obtained are shown in Table 1 below.

Table 1

Tested Compound	PKC-IC ₅₀ (μM)	PKA-IC ₅₀ (μM)	PKA-IC ₅₀ /PKC-IC ₅₀
Polymyxin B Sulfate	10	>100	>10
Calphostin C	0.05	>50	>1000
Palmitoyl-DL-carnitine Hydrochloride	100	>300	>3
hexadecylphosphocholine	94	>1000	>10.6
H-7 ^{*1}	15	13	0.87
K252a ^{*1}	0.02	0.02	1
Staurosporine ^{*1}	0.0027	0.0054	2

*1: Excerpt from BIO/TECHNOLOGY, 8, 732, 1990.

Reference Example 2: Preparation of Tonic 5 to 9 containing Staurosporine

To the mixed solution containing 90 g of ethyl alcohol, 5 g of 1,3-butyleneglycol, 0.5 g of N-acetylglutamine iso-stearyl ester and 0.25 g of polyoxyethylene(25) glyceryl pyroglutamate isostearate diester added 1×10^{-6} g, 3×10^{-6} g, 1×10^{-5} g or 3×10^{-5} g, 1×10^{-4} g of Staurosporine, and was uniformly mixed and stirred to prepare 5 kinds of solution.

To these solutions was added 4.22 g of pure water respectively with stirring, and uniformly mixed to prepare hair growing tonics containing 0.02 μM (composition 5), 0.06 μM (composition 6), 0.2 μM (composition 7), 0.6 μM (composition 8), 2 μM (composition 9) of Staurosporine.

Test Example 1: Hair Growth Promoting Effect in Mouse Hair-Follicle Cell Culture

Mouse hair-follicle cells were cultivated and collected, according to the modified method of the Tanigai et al's method (see Archives of Dermatological Research, 284, 290-296, 1992) while suitably modifying the method.

That is, the skin was peeled from the back of a 4-day age C3H mouse (bought from Nippon Charles River Co.), and processed with an MEM (Minimum Essential Medium, Eagle) containing 500 units/ml Disperse (produced by Godo Shusei Co.) and 5 % FCS, at 4°C for 16 hours.

The epidermis was removed from said skin sample, and the resulting dermis layer was processed with a DMEM (Dulbecco's Modified Eagle Medium) containing 0.25 % collagenase N-2 (produced by Nitta Gelatin Co.) and 10 % FCS, at 37°C for 1 hours, to prepare a dermis suspension.

This dermis suspension was filtrated through a 212-micron nylon mesh (produced by Nippon Rikagaku Kikai KK), and the resulting filtrate was centrifuged at 1,000 rpm for 5 minutes to obtain hair-follicle tissue-containing pellets.

The resulting pellets were suspended in a Ca-Mg-free PBS (Dulbecco's Phosphate-Buffered Saline), using a pipette, and then statically left as it was for 15 minutes, whereby the hair-follicle tissue was precipitated.

The thus-obtained hair-follicle tissue was repeatedly processed for a total of three times according to the same process as above comprising the addition of a Ca-Mg-free PBS, the suspension with a pipette, and the precipitation for 15 minutes.

The hair-follicle tissue thus obtained finally was processed with 0.1 % EDTA-0.25 % trypsin (produced by Gibco Co.) at 37°C for 5 minutes, to which was then added a DMEM containing 10 % FCS. Thus was prepared herein a hair-follicle tissue cell suspension comprising 3×10^5 cells/ml.

This hair-follicle tissue cell suspension was pipetted into a 24-well collagen-coated plate (produced by Iwaki Glass Co.) in an amount of 1 ml/well, and cultivated at 37°C in 5 % CO₂ for 24 hours.

The resulting culture was then subjected to medium exchange with an MCDB153 medium (produced by Kyokuto Pharmaceutical Co.) containing 5 mg/liter insulin from bovine (produced by Sigma Co.), 5 µg/liter mouse EGF (produced by Takara Shuzo Co.), 40 mg/liter bovine pituitary extract (produced by Kyokuto Pharmaceutical Co.), 10 mg/liter human transferrin (produced by Sigma Co.), 0.4 mg/liter hydrocortisone (produced by Sigma Co.), 0.63 µg/liter progesterone (produced by Collaborative Research Co.), 14 mg/liter O-phosphoethanolamine (produced by Sigma Co.), 6.1 mg/liter ethanolamine (produced by Sigma Co.), 50 U/ml penicillin (produced by Wako Co.), 50 µg/ml streptomycin (produced by Wako Co.), and an DMSO solution comprising a PKC inhibitor of the invention (this was added in an amount of 1/100 by volume relative to the medium), and then further cultivated at 37°C in 5 % CO₂ for 5 days. During the incubation, the medium was exchanged with a fresh one every other day.

As the control, the cell culture was cultivated in the same manner as above, except that only DMSO of 1/100 by volume relative to the medium was added to the medium in place of the PKC inhibitor-containing DMSO solution.

To determine the degree of cell growth, referred to herein was a known method of using Neutral Red (see Journal of Tissue Culture Method, 9, 1, 7-9, 1984).

The resulting culture was extracted through suction, and then further cultivated in an MCDB153 medium, to which had been added 50 mg/liter Neutral Red (produced by Sigma Co.), at 37°C in 5 % CO₂ for 3 hours. The supernatant was removed from the culture, and the remaining cells were washed with a 1 % formalin solution containing 1 % calcium chloride, and then fixed.

After having been thus washed and fixed, the supernatant was removed, and a 50 % ethanol solution containing 1 % acetic acid was added to the residue in an amount of 0.4 ml/well of 2 cm², whereby Neutral Red was extracted out.

The absorbance of the extract at 540 nm was measured, from which was obtained the degree of growth of the cells.

The cell growth-promoting activity of each tested compound of the invention tested herein is shown in Table 2 below. As in Table 2, the hair-growing agent comprising each compound of the invention exhibits an excellent hair-growing effect of the mouse hair-follicle cell.

On the other hand, protein kinase inhibitors, K252a and Staurosporine, having PKA-IC₅₀/PKC-IC₅₀ of smaller than 3 were tested in the same manner as above, and the data obtained are shown in Table 3 below. As in Table 3, these protein kinase inhibitors inhibited the growth of mouse hair-follicle cells.

Table 2

Compound Tested	Concentration (µM)	Relative Degree of Cell Growth, vehicled on the degree of cell growth of 100 in Control
Polymyxin B Sulfate	10	360
Calphostin C	0.1	160
Palmitoyl-DL-carnitine Hydrochloride	10	170

Table 3

Compound Tested	Concentration (μ M)	Relative Degree of Cell Growth, vehicled on the degree of cell growth of 100 in Control
K252a	1×10^5	80
	1×10^4	77
	1×10^3	76
	1×10^2	29
	1×10^1	0
Staurosporine	1×10^5	78
	1×10^4	69
	1×10^3	72
	1×10^2	34
	1×10^1	0

Test Example 2: Effect on Mouse Hair Growth

The compositions prepared hereinabove were tested for their effects on the growth of hair of mice, with reference to the Ogawa et al's method (see The Journal of Dermatology, Vol. 10, pp. 45-54, 1983).

Precisely, 9-week age, male C3H/HeSlc mice, of which the hair cycle was in the resting period, were grouped into plural groups each comprised of 4 or 5 mice. The hair on the back of each mouse was carefully shaven, using electric hair clippers and an electric shaver, and any of the compositions 1 to 9 as prepared in Examples 1 to 4 and Reference Example 2 was uniformly applied to the shaven part of each mouse, once a day in an amount of 200 μ l/part.

To the mice of the control group, applied was the same composition as above except that it did not contain the protein kinase inhibitor of the invention.

On the 18th day after the start of the application of the composition to mice, the shaven skin on the back of each mouse was peeled and photographed. The photographic pictures obtained were imaged, using an image processor (produced by Avionics Co., Spicca II), whereby the percentage of the hair-grown area relative to the total area of the shaven skin was obtained to be the degree of hair growth. The value, by which the value of hair-growing percentage of the group to which the chemical had been applied is subtracted from the value of hair-growing percentage of the control group, was obtained to be the percentage of area with increased hair growth (%).

The data obtained herein are shown in Table 4 and Table 5 below.

As shown in Table 4, the hair-growing agents comprising the protein kinase C-specific inhibitor of the invention (composition 1 to 4) exhibit an excellent hair-growing effect of the mouse hair-follicle.

On the other hand, as shown in Table 5, no or little such effect was exhibited by using composition 5 to 9 containing Staurosporine which is protein kinase inhibitor having PKA-IC50/PKC-IC50 of smaller than 3 (composition 5 to 9).

Table 4

Test Composition	Percentage of Area with Increased Hair Growth (%)
Composition 1	58
Composition 2	65
Composition 3	51
Composition 4	70

Table 5

Test Composition	Concentration of Staurosporine(μ M)	Percentage of Area with Increased Hair Growth (%)
Composition 5	0.02	-2
Composition 6	0.06	-5
Composition 7	0.20	3
Composition 8	0.60	-8
Composition 9	2.00	-14

As has been mentioned in detail hereinabove, the present invention provides a safe and effective hair-growing agent comprising a protein kinase C-specific inhibitor.

While the invention has been described in detail and with reference to specific embodiments thereof, it will be apparent to one skilled in the art that various changes and modifications can be made therein without departing from the spirit and scope thereof.

Claims

1. A hair-growing agent comprising, as an active ingredient, a protein kinase C-specific inhibitor and pharmaceutically acceptable vehicles.
2. The hair-growing agent according to claim 1, wherein the protein kinase C-specific inhibitor is a protein kinase inhibitor in which a ratio of its 50 % protein kinase A-inhibiting constant to its 50 % protein kinase C-inhibiting constant is not smaller than 3.0.
3. The hair-growing agent according to claim 1, wherein the protein kinase C-specific inhibitor is a protein kinase inhibitor in which a ratio of its 50 % protein kinase A-inhibiting constant to its 50 % protein kinase C-inhibiting constant is from 3.0 to 10^9 .
4. The hair-growing agent according to claim 1, wherein the protein kinase C-specific inhibitor is a protein kinase inhibitor in which a ratio of its 50 % protein kinase A-inhibiting constant to its 50 % protein kinase C-inhibiting constant of from 10 to 10^9 .
5. The hair-growing agent according to claim 1, 2, 3 or 4, wherein the protein kinase C-specific inhibitor is selected from polymyxin B, calphostin C, palmitoyl-DL-carnitine, hexadecylphosphocholine and their pharmaceutically-acceptable salts.
6. The hair-growing agent according to claim 1, wherein the amount of the protein kinase C-specific inhibitor to be contained in the agent is 10^{-6} to 10 % by weight based on the agent.